

PLURIPOLARITY OF GRAPHS OF QUASIANALYTIC FUNCTIONS IN THE SENSE OF GONCHAR

SEVDIYOR IMOMKULOV AND ZAFAR IBRAGIMOV

Let f be a continuous real valued function, defined on a segment $\Delta = [a, b]$ of the real axis $\mathbb{R}$ and let $\rho_n(f)$ be the deviation from the best approximation of f on Δ by rational functions $r_n(x)$ of degree less than or equal to n . That is,

$$\rho_n(f) = \inf_{r_n} \max_{x \in \Delta} |f(x) - r_n(x)|,$$

where the infimum is taken over all rational functions of the form

$$r_n(x) = \frac{a_0 x^n + a_1 x^{n-1} + \dots + a_n}{b_0 x^n + b_1 x^{n-1} + \dots + b_n}.$$

As usual, we denote by $e_n(f)$ the least deviation of f on Δ from its polynomial approximation of degree less than or equal to n . Obvious, $\rho_n(f) \leq e_n(f)$ for every $n = 0, 1, 2, \dots$. In papers ([1],[2]) Gonchar proved that the class of functions

$$R(\Delta) = \{f \in C(\Delta) : \lim_{n \rightarrow \infty} \sqrt[n]{\rho_n(f)} < 1\}$$

possesses one of the important properties of the class of analytic functions. Namely, if

$$\lim_{n \rightarrow \infty} \sqrt[n]{\rho_n(f)} < 1$$

and $f(x) = 0$ on a set $E \subset \Delta$ of positive logarithmic capacity, then $f(x) \equiv 0$ on Δ .

By analogy with the class

$$B(\Delta) = \{f \in C(\Delta) : \lim_{n \rightarrow \infty} \sqrt[n]{e_n(f)} < 1\},$$

which is called the class of quasianalytic functions of Bernstein (see [3]), we call $R(\Delta)$ the class of quasianalytic functions of Gonchar.

It is known that functions that are analytic on Δ are characterized by condition

$$\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{e_n(f)} < 1,$$

according to Bernstein's theorem. It follows that the class $A(\Delta)$ of analytic functions on Δ is a subclass of $B(\Delta) \subset R(\Delta)$, i.e., $A(\Delta) \subset B(\Delta) \subset R(\Delta)$. It is clear that $A(\Delta) \neq B(\Delta)$. It was shown in ([1]) that there exist functions for which the rate of approximation by polynomials $e_n(f)$ tends to zero as slow as possible whereas $\rho_n(f)$ tends to zero as fast as possible. In particular, it follows that

$$B(\Delta) \neq R(\Delta).$$

It is not hard to see that if $f \in A(\Delta)$, then its graph

$$\Gamma_f = \{(x, f(x)) \in \mathbb{C}^2 : x \in \Delta\}$$

is a pluripolar set in $\mathbb{C}^2$. In ([5]) the authors constructed examples of smooth (infinitely differentiable) functions whose graph are not pluripolar in $\mathbb{C}^2$. In their recent paper ([3]) the authors have proved that if $f \in B(\Delta)$, then its graph Γ_f is pluripolar in $\mathbb{C}^2$.

In this paper we prove following more general theorem.

Theorem 1. *If $f \in R(\Delta)$, then its graph Γ_f is pluripolar in $\mathbb{C}^2$.*

Proof. By the hypothesis of the theorem there exists a sequence of natural numbers n_k and a corresponding sequence of rational functions

$$r_{n_k} = \frac{p_{n_k}}{q_{n_k}}$$

such that

$$\rho_{n_k}(f) = \left\| f - r_{n_k} \right\|_{\Delta} \leq \alpha^{n_k},$$

where $\alpha \in (0, 1)$ is some fixed number. Without loss of generality we can assume that

$$\left\| f \right\|_{\Delta} \leq \frac{1}{2}, \quad \left\| p_{n_k} \right\|_{\Delta} \leq 1 \quad \text{and} \quad \left\| q_{n_k} \right\|_{\Delta} = 1.$$

According to Bernstein-Walsh inequality (see, [7]) we have

$$\left| p_{n_k}(z) \right| \leq e^{n_k V^*(z, \Delta)} \quad \text{and} \quad \left| q_{n_k}(z) \right| \leq e^{n_k V^*(z, \Delta)}$$

for any $z \in \mathbb{C}$ and $k \in N$. Here

$$V^*(z, \Delta) = \overline{\lim}_{z' \rightarrow z} \sup \left\{ \frac{1}{n} \ln |p_n(z')| : \|p_n\|_{\Delta} \leq 1 \right\}$$

is the extremal function of Green.

We introduce the following auxiliary sequence of plurisubharmonic functions

$$u_k(z, w) = \frac{1}{n_k} \ln |q_{n_k}(z) \cdot w - p_{n_k}(z)|, \quad (z, w) \in \mathbb{C}^2.$$

For $(z, w) \in \mathbb{C}^2$ we have

$$\begin{aligned} \frac{1}{n_k} \ln |q_{n_k}(z) \cdot w - p_{n_k}(z)| &\leq \frac{1}{n_k} \ln (|q_{n_k}(z) \cdot w| + |p_{n_k}(z)|) \\ &\leq \max \left\{ \frac{1}{n_k} \ln 2 |p_k(z)|, \frac{1}{n_k} \ln 2 |q_{n_k}(z) \cdot w| \right\} \\ &= \max \left\{ \frac{1}{n_k} \ln |p_k(z)|, \frac{1}{n_k} \ln |q_{n_k}(z)| + \frac{1}{n_k} \ln |w| \right\} + \frac{\ln 2}{n_k}. \end{aligned}$$

From here we obtain the following estimate

$$u_k(z, w) \leq \max \{ V^*(z, \Delta), V^*(z, \Delta) + \frac{1}{n_k} \ln |w| \} + \frac{\ln 2}{n_k}.$$

Consequently, the sequence of plurisubharmonic functions $u_k(z, w)$ is locally uniformly bounded from above. Let

$$u(z, w) = \overline{\lim}_{k \rightarrow \infty} u_k(z, w).$$

The function $u(z, w)$ is also locally bounded from above,

$$u(z, w) \leq V^*(z, \Delta).$$

Let

$$u^*(z, w) = \overline{\lim}_{(z', w') \rightarrow (z, w)} u(z', w')$$

denote the regularization of function $u(z, w)$. The set

$$E = \{(z, w) \in \mathbb{C}^2 : u(z, w) < u^*(z, w)\}$$

is pluripolar in $\mathbb{C}^2$ (see [6],[7]).

Let now $(z, w) \in \Gamma_f$ be a fixed point. (Note that $q_n(z) \neq 0$ for $z \in \Delta$). Then

$$\begin{aligned} u(z, w) &= \overline{\lim}_{k \rightarrow \infty} \ln |q_{n_k}(z)|^{\frac{1}{n_k}} \left| w - \frac{p_{n_k}(z)}{q_{n_k}(z)} \right|^{\frac{1}{n_k}} \leq \overline{\lim}_{k \rightarrow \infty} \ln \alpha |q_{n_k}(z)|^{\frac{1}{n_k}} \\ &= \ln \alpha + \overline{\lim}_{k \rightarrow \infty} \ln |q_{n_k}(z)|^{\frac{1}{n_k}}. \end{aligned}$$

If $(z, w) \in (\Delta \times \mathbb{C}) \setminus \Gamma_f$, then

$$\begin{aligned} u(z, w) &= \overline{\lim}_{k \rightarrow \infty} \frac{1}{n_k} \ln |q_{n_k}(z)w - p_{n_k}(z)| \\ &= \overline{\lim}_{k \rightarrow \infty} \ln |q_{n_k}(z)|^{\frac{1}{n_k}} \left| w - \frac{p_{n_k}(z)}{q_{n_k}(z)} \right|^{\frac{1}{n_k}} = \overline{\lim}_{k \rightarrow \infty} \ln |q_{n_k}(z)|^{\frac{1}{n_k}}. \end{aligned}$$

It follows that if

$$\overline{\lim}_{k \rightarrow \infty} |q_{n_k}(z)|^{\frac{1}{n_k}} \neq 0$$

at a point $z \in \Delta$, then $(z, f(z))$ belongs to the pluripolar set E . Therefore, to complete the proof of the theorem it is enough to show that the set

$$A = \left\{ z \in \Delta : \overline{\lim}_{k \rightarrow \infty} |q_{n_k}(z)|^{\frac{1}{n_k}} = 0 \right\} = \left\{ z \in \Delta : \lim_{k \rightarrow \infty} |q_{n_k}(z)|^{\frac{1}{n_k}} = 0 \right\}$$

is polar. Assume that A is not polar, i.e., it has a positive logarithmic capacity, $\text{cap}(A) > 0$.

We consider the following sequence of subharmonic functions

$$\vartheta_k^*(z) = \overline{\lim}_{z' \rightarrow z} \vartheta_k(z'), \quad z \in \mathbb{C},$$

where

$$\vartheta_k(z) = \sup_{m \geq k} |q_{n_m}(z)|^{\frac{1}{n_m}}.$$

It is clear that the sequence $\vartheta_k^*(z)$ is locally uniformly bounded, $0 \leq \vartheta_k^*(z) \leq e^{V^*(z, \Delta)}$ and is not monotonically increasing. In addition, $\vartheta_k^*(z) \rightarrow 0$ on A , except for the polar set

$$F = \bigcup_{k=1}^{\infty} \{z \in \mathbb{C} : \vartheta_k(z) < \vartheta_k^*(z)\}$$

since by definition the sequence $\vartheta_k(z)$ tends to zero on A .

Since $\vartheta_k^*(z)$ is monotonic, for every ϵ , $0 < \epsilon < \text{cap}(A)$ there exists an open set U_ϵ , $\text{cap}(U_\epsilon) < \epsilon$ such that the sequence $\vartheta_k^*(z)$ converges uniformly on the

set $A_\epsilon = A \setminus U_\epsilon$ (see, for example, [6]). It follows that there exists a compact set $A_0 \subset A_\epsilon$, $\text{cap}(A_0) > 0$, such that the sequence of subharmonic functions $\vartheta_k^*(z)$ converges uniformly to zero on the set A_0 . Consequently, the sequence $|q_{n_k}(z)|^{1/n_k}$ also converges uniformly to zero on the compact set A_0 . Now we use the so-called τ -capacity, introduced by A.Sadullaev (see [6]). We consider a polynomial $T_n(z)$, $\deg T_n(z) \leq n$, normalized by condition $\|T_n\|_{U(0,r)} = 1$, where $U(0,r) = \{z: |z| < r\} \supset \Delta$ for which the norm $\|T_n\|_{A_0}$ is minimal among all such polynomials. Then the limit

$$\lim_{n \rightarrow \infty} \|T_n\|_{A_0}^{\frac{1}{n}} = \tau(A_0, U(0, r)),$$

exists and, moreover

$$\|T_n\|_{A_0}^{\frac{1}{n}} \geq \tau(A_0, U(0, r))$$

for every $n \in N$. In fact,

$$\tau(A_0, U(0, r)) = \exp\left\{-\max_{z \in U(0,r)} V^*(z, A_0)\right\}.$$

It follows that

$$(0.1) \quad \|q_{n_k}\|_{A_0}^{\frac{1}{n_k}} \geq \|T_{n_k}\|_{A_0}^{\frac{1}{n_k}} \geq \tau(A_0, U(0, r)) > 0$$

for

$$\|q_{n_k}\|_{\Delta} = 1 \quad \text{and} \quad \|q_{n_k}\|_{U(0,r)} \geq 1.$$

It follows from inequality (0.1) that the sequence $|q_{n_k}(z)|^{\frac{1}{n_k}}$ does not converge uniformly to zero on the set A_0 . We arrived at a contradiction. Hence the set A is polar.

Thus, the graph Γ_f is divided into two parts, $\Gamma_f = \Gamma'_f \cup \Gamma''_f$, where

$$\Gamma'_f = \{(z, f(z)): z \in \Delta \setminus A\} \quad \text{and} \quad \Gamma''_f = \{(z, f(z)): z \in A\}.$$

The first set is a subset of a pluripolar set $E \subset \mathbb{C}^2$ and the second set is a subset of a pluripolar set $A \times \mathbb{C} \subset \mathbb{C}^2$. Consequently, Γ_f is a pluripolar in $\mathbb{C}^2$. The proof of the theorem is complete. □

Sadullaev's additions. Actually, in this work the authors proved a stronger result. Let K be arbitrary compact set in $\mathbb{C}^n$ and let $f(z) \in C(K)$. We say that $f(z)$ belongs to the class of Gonchar $R(K)$ if

$$\liminf_{k \rightarrow \infty} \rho_k^{1/k}(f, K) < 1, \quad \text{where} \quad \rho_k(f, K) = \inf_{\{r_k\}} \max_{z \in K} |f(z) - r_k(z)|$$

and the infimum is taken over all rational functions $r_k(z)$ of degree less than or equal to k . The following more general result holds: if the function $f(z)$ belongs to the class of Gonchar $R(K)$, then its graph $\Gamma_f = \{(z, f(z)) \in \mathbb{C}^{n+1} : z \in K\}$ is pluripolar in $\mathbb{C}^{n+1}$.

We note that pluripolarity of a graph Γ_f is closely related to analyticity property of $f(z)$ or to properties of $f(z)$ related to analyticity. Analyticity of continuous functions in domains $D \subset \mathbb{C}^n$ whose graphs are pluripolar in $\mathbb{C}^{n+1}$ was proved in ([10]). An example of a function $f(z)$, which is holomorphic in the unit disk $U \subset \mathbb{C}$ and continuous in the closed disk $\overline{U}$, and whose graph Γ_f ,

$$\Gamma_f = \{(z, f(z)) \in \mathbb{C}^2 : z \in \overline{U}\},$$

is not pluripolar is constructed in ([9]). These results suggest that the main reason behind pluripolarity of graphs is the analyticity of functions.

In ([8]) an example of a lacunary series

$$f(z) = \sum_{k=1}^{\infty} a_{n_k} z^{n_k} \quad \left(\lim_{k \rightarrow \infty} \sqrt[n_k]{a_{n_k}} = 1, \quad \frac{a_{n_k}}{a_{n_k+1}} \rightarrow 0 \quad \text{as } k \rightarrow \infty \right)$$

whose graph Γ_f is plurispaced in every boundary point. It would be interesting to establish if the graph of a function $f(z)$,

$$f(z) = \sum_{k=1}^{\infty} \frac{1}{k^{\ln k}} z^{k!},$$

which is infinitely smooth up to the closure of its domain, is pluripolar in $\mathbb{C}^2$? If we consider the series

$$f(z) = \sum_{k=1}^{\infty} \frac{1}{2^{(k-1)!}} z^{k!},$$

then its graph Γ_f will be pluripolar since $f(z)$ is quasianalytic in the sense of Gonchar.

REFERENCES

- [1] Gonchar A. A., *On best approximations by rational functions*, Dokl. Akad. Nauk SSSR, 100 (1955), 205–208, (Russian).
- [2] Gonchar A. A., *Quasianalytic class of functions, connected with best approximated by rational functions*, Izv. of the Academy of Sciences Armenia SSR. 1971, VI . No. 2-3, 148–159.
- [3] Coman, D., Levenberg, N. and Poletsky, E. A., *Quasianalyticity and Pluripolarity*, arxiv:math/0402381.v1[math.CV] 23 feb, 2004.
- [4] Timan, A. F., *Theory of approximation of functions of a real variable*, Pergamon Press, Macmillan, New York, 1963.
- [5] Diederich, K. and Fornass, J. E., *A smooth curve which is not a pluripolar set*, Duke Math.J. 49 (1982), 931–936.
- [6] Sadullaev, A., *Rational approximation and pluripolar sets*, Mat. Sb., 119:1 (1982), 96–118; English transl. in Math. USSR-Sb., 47 (1984).
- [7] Sadullaev, A., *Plurisubharmonic functions*, Sovrem. Probl. Mat. Fund. Naprav, 8, (VINITI, Moscow), 1985, 65–113 ; English transl. in Encyclopaedia Math. Sci., vol. 8 [Several complex variables II], (Springer-Verlag, Berlin), 1984.
- [8] Sadullaev, A., *Plurisubharmonic measures and capacities on complex manifolds*, Uspekhi Mat. Nauk, 36:4 (1981), 53–105; English transl. in Russian Math. Surveys, 36 (1981).
- [9] Levenberg, N., Martin, G. and Poletskiy, E. A., *Analytic disks and pluripolar sets*, Indiana Univ. Math. J. 41 (1992), 515–532.
- [10] Shcherbina, N., *Pluripolar graphs are holomorphic*, Acta Math., 194 (2005), 203–216.